




International Journal of Alternative and Complementary Medicine

Available at www.saapjournals.org

ISSN: 2583-1305



HOMEOPATHY IN CHRONIC DISEASE MANAGEMENT: A CRITICAL REVIEW OF THE EVIDENCE

Tithy Rani Sharkar¹ and Abdul Gaffar Miah*²¹Department of Materia Medica, Natore Homeo Medical College, Natore, Bangladesh.²Department of Genetics and Animal Breeding, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh.

Received: 22 Aug 2025 Revised: 11 Sept 2025 Accepted: 16 Nov 2025

Abstract

Homeopathy remains one of the most widely practiced complementary and alternative medicine (CAM) modalities worldwide, particularly among individuals with chronic non-communicable diseases (NCDs) who seek safe, holistic, and personalized therapeutic options. Despite its global popularity, controversy continues regarding its mechanisms of action and clinical effectiveness. This review critically evaluates the current evidence on the role of homeopathy in chronic disease management. A comprehensive review of randomized controlled trials (RCTs), systematic reviews, and meta-analyses published between 2000 and 2025 was conducted. Studies were included if they examined homeopathic interventions in chronic conditions such as asthma, arthritis, type 2 diabetes mellitus, depression, chronic pain, and fibromyalgia. Data were synthesized thematically to assess efficacy, mechanistic plausibility, methodological quality, and safety outcomes. Mechanistic hypotheses suggest that hormesis, nanoparticle-mediated signaling, immune modulation, and neuropsychological or psychosocial mechanisms may contribute to the therapeutic effects of homeopathy. Meta-analytic evidence demonstrates modest but statistically significant improvements in subjective measures such as pain, fatigue, and quality of life across several chronic diseases, with standardized mean differences ranging from 0.18 to 0.25. However, objective clinical outcomes, including spirometry and HbA1c, rarely show consistent benefit. Methodological challenges, including small sample sizes, heterogeneous interventions, limited follow-up durations, and a high risk of bias, continue to constrain the reliability of existing findings. Importantly, homeopathy exhibits a favorable safety profile, with no serious adverse events reported across chronic disease studies. Current evidence indicates that homeopathy may provide modest adjunctive benefits for symptom relief and improved patient satisfaction among individuals with chronic illnesses, although its specific therapeutic mechanisms remain uncertain. Integration of homeopathy into evidence-based, patient-centered chronic disease management frameworks may enhance holistic care. Future research should emphasize large-scale, multicenter randomized trials with standardized outcome measures and mechanistic endpoints to better define clinical relevance and biological plausibility.

Keywords: Homeopathy; Chronic disease management; Complementary and alternative medicine; Meta analysis; Immunomodulation

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. Copyright © 2025 Author(s) retain the copyright of this article.



*Corresponding Author

Professor Dr. Abdul Gaffar Miah

DOI: <https://doi.org/10.46797/ijacm.v6i2.777>

Produced and Published by

South Asian Academic Publications

Introduction

Chronic non-communicable diseases (NCDs) such as cardiovascular diseases, cancer, diabetes and chronic respiratory diseases remain the leading causes of global mortality and disability. According to the World Health Organization (WHO), NCDs were responsible for

approximately 74 percent of global deaths in 2019, representing about 41 million deaths worldwide [1]. Cardiovascular diseases accounted for 17.9 million deaths, cancer for 9.3 million, chronic respiratory diseases for 4.1 million and diabetes for 1.5 million [1]. Although age-specific mortality rates have declined in many regions, the total number of individuals living with NCDs continues to increase due to population ageing and overall population expansion [1]. This rising burden contributes to premature mortality and creates substantial economic and social challenges for countries globally [2].

Management of chronic diseases typically requires long-term medication, regular monitoring and sustained lifestyle modifications. Although these interventions are

effective, many patients experience side effects, treatment burden and low adherence. As a result, a significant proportion of patients seek complementary and alternative medicine (CAM) as either an adjunct or an alternative to conventional treatment. CAM includes a wide range of practices such as herbal medicine, acupuncture, mindfulness therapy and homeopathy that are generally not integrated within conventional biomedical systems [3]. Among these, homeopathy remains widely used, particularly by individuals with chronic diseases who do not achieve satisfactory relief from standard treatments [4].

Homeopathy was developed by Samuel Hahnemann in the late eighteenth century and is based on the principle of "like cures like," meaning that substances capable of producing symptoms in healthy individuals are used in highly diluted forms to treat similar symptoms in patients. Homeopathic remedies are prepared through serial dilution with succussion, a method known as potentization. Despite ongoing debate regarding the scientific plausibility of such high dilutions, homeopathy continues to be used extensively across many regions. Surveys in Europe and parts of Asia report homeopathy use among 20 to 40 percent of patients with chronic health conditions [4,5]. In a multicentre study conducted in Bangladesh, 32.8 percent of adults with chronic illnesses reported using CAM, and 52.2 percent of these individuals used homeopathy [5].

The continued popularity of homeopathy, despite longstanding scientific controversy, highlights the need for an updated and critical appraisal of its role in chronic disease management. Several randomized controlled trials (RCTs) and systematic reviews have evaluated homeopathic interventions in asthma, osteoarthritis, diabetes, chronic pain and other long-term conditions, although findings are often mixed and limited by methodological weaknesses such as small sample sizes, heterogeneity of interventions and a high risk of bias [4,6]. Proposed mechanisms include hormesis, nanoparticle-related activity, immunomodulation and placebo-associated neuropsychological pathways, although these mechanisms require stronger experimental validation [6].

The primary aim of this review is to synthesise and critically evaluate the current evidence on homeopathic interventions in chronic disease management, drawing on RCTs, observational studies, systematic reviews and meta-analyses. The specific objectives are to (a) assess efficacy and safety outcomes, (b) examine proposed mechanisms and their plausibility, (c) identify methodological strengths and limitations and (d) provide evidence-informed guidance for clinicians, researchers and policymakers. Given the long-term nature of chronic diseases and the significant impact of treatment decisions on quality of life, health outcomes and economic burden, evidence-based assessment of homeopathy is essential to support safe and effective patient care [2,3].

Historical and Theoretical Background of Homeopathy (Expanded)

Homeopathy originated in the late eighteenth century through the work of Samuel Hahnemann, a German physician who sought alternatives to the harmful medical practices of his time, including bloodletting and the use of toxic compounds such as mercury [7]. Hahnemann proposed the principle of *similia similibus curentur* or "like cures like," which suggests that substances producing certain symptoms in healthy individuals can be used in highly diluted forms to treat similar symptoms in diseased individuals. This principle remains central to homeopathic practice.

Hahnemann also emphasized individualized treatment, arguing that therapy should be based on the totality of a patient's symptoms rather than a single disease label. This individualized approach continues to distinguish homeopathy from many other complementary or conventional medical systems [7, 8].

Homeopathic remedies are prepared through serial dilution combined with vigorous shaking, a process termed potentization. Hahnemann hypothesized that this process released a form of medicinal energy from the original substance that could stimulate the body's self-healing capacity [7]. Contemporary homeopathy often asserts that potentization both reduces toxicity and enhances therapeutic potential. Critics, however, argue that at high dilutions commonly used in practice, remedies may contain no detectable molecules of the original substance, raising concerns about pharmacological plausibility [9, 10].

Homeopathy gained rapid popularity during the nineteenth century in Europe and North America, partly due to dissatisfaction with conventional medicine of that era. Homeopathic institutions, hospitals and training centres proliferated, contributing to its recognition and use among the public. With the rise of scientific medicine in the early twentieth century, including advances in pharmacology and microbiology, homeopathy's prominence decreased within mainstream healthcare. Nevertheless, it continued to be practiced in various regions and later experienced renewed global interest as patients increasingly sought holistic and less invasive approaches for chronic disease management [7, 8].

A number of hypotheses have been advanced to explain the potential biological effects of homeopathic remedies. One proposed mechanism is hormesis, in which low doses of bioactive substances may stimulate adaptive cellular responses with potential therapeutic benefits [11]. In vitro experiments show that low concentrations of certain plant-derived compounds can influence antioxidant enzyme activity, gene expression and cellular stress responses [7, 8]. Such findings may support the use of homeopathy in chronic conditions involving chronic inflammation and oxidative stress.

Another proposed mechanism is the nanoparticle hypothesis. Studies indicate that even at high dilutions,

homeopathic preparations may contain nanoparticles or nanostructures derived from the original substance [12]. These structures may interact with biological systems and influence signalling pathways, although evidence remains preliminary. Related to this idea is the water structure or water memory hypothesis, which suggests that the solvent retains a structural imprint of the original substance. This hypothesis remains controversial, as findings have been inconsistent across independent studies [7].

Immunomodulation has also been proposed as a potential mechanism. Preclinical studies report that homeopathic preparations may influence cytokine production, inflammatory signalling and immune cell activity [7, 8]. In animal models of asthma and arthritis, some homeopathic substances have been shown to modulate inflammatory mediators such as TNF alpha, IL 6 and IL 10 [8, 13]. Although these findings suggest possible biological effects, reproducibility and translation into clinical outcomes in humans remain uncertain.

Debate about homeopathy’s mechanisms persists. Critics point out that many mechanistic studies suffer from small sample sizes, lack of replication and methodological variability [9,10]. Others argue that placebo and contextual effects account for much of the perceived benefit of homeopathy, especially in conditions where subjective symptoms dominate [10]. Supporters of homeopathy contend that its individualized care, minimal side effects and patient satisfaction justify its use, particularly when conventional treatments are inadequate or poorly tolerated.

Homeopathy’s continued use is reflected in global CAM utilisation patterns. Surveys from Bangladesh and Europe indicate substantial use of homeopathy among individuals with chronic diseases, including asthma, arthritis, diabetes and chronic pain [5,8]. In Bangladesh, 32.8 percent of adults with chronic illnesses reported using CAM, with homeopathy being the most frequently used modality [5].

In summary, homeopathy integrates historical principles of individualized care with several theoretical mechanisms that remain debated in contemporary science. Hormesis, nanoparticle effects and immunomodulation have been proposed, but evidence remains limited and inconsistent. Despite these controversies, homeopathy continues to play a role in chronic disease care, highlighting the need for systematic evaluation of its mechanisms and clinical effectiveness. The following sections provide critical assessment of mechanistic hypotheses and clinical evidence across chronic disease categories.

Proposed Mechanisms of Action of Homeopathy

The mechanisms underlying homeopathic effects remain a topic of ongoing debate in modern biomedical science. The extremely high dilutions used in many homeopathic preparations challenge conventional pharmacological principles and contribute to controversy regarding biological plausibility [9, 10]. Nevertheless, several hypotheses have been proposed to explain potential

therapeutic effects. These can be grouped into low dose biological responses, nanoparticle related hypotheses, immunomodulatory pathways, and neuropsychological and psychosocial mechanisms.

1. Hormesis and Low Dose Biological Stimulation

Hormesis refers to a biphasic dose response relationship in which low doses of a biologically active substance can stimulate beneficial adaptive responses, whereas high doses may be inhibitory or toxic [11]. In homeopathy, it is proposed that highly diluted substances may function as hormetic triggers that induce mild cellular stress, thereby activating homeostatic and adaptive pathways. Experimental studies have shown that low concentrations of plant derived compounds can influence antioxidant enzyme activity, gene expression and cellular stress resistance [7, 8]. These findings provide theoretical support for potential benefits in chronic conditions that involve oxidative stress and inflammation, such as asthma and arthritis. However, robust clinical validation of hormetic mechanisms in human populations receiving homeopathic treatment remains limited.

A range of biological and theoretical models has been proposed to explain how highly diluted homeopathic preparations might exert physiological effects. These models include hormesis, nanoparticle related activity, solvent structure changes and immunomodulatory pathways, each supported to varying extents by preclinical or clinical observations [7,11-13]. To provide a concise overview of these concepts, the principal mechanistic hypotheses and their potential relevance to chronic disease are summarised in Table 01. This table highlights the proposed pathways, key supporting evidence and major limitations, offering a foundation for interpreting subsequent sections of this review.

Table 01: Proposed mechanistic hypotheses of homeopathy in chronic diseases.

Mechanism	Proposed Effect	Evidence	References
Hormesis / Ultra-low dose effect	Activation of adaptive cellular responses	<i>In vitro</i> and preclinical studies	Bellavite et al., 2021
Nanoparticle / water-structure hypothesis	Modulation of biochemical signaling pathways	Experimental and theoretical studies	Jacobs et al., 2022
Immunomodulation	Alteration of cytokines (IL-6, TNF-α)	Animal models and some clinical trials	Bellavite et al., 2021; Patwardhan et al., 2021
Neuropsychology	Modulation	Clinical	Kaptchuk

ical / placebo	n of brain reward and stress pathways	trials, neuroimaging studies	et al., 2020
Psychosocial / therapeutic encounter	Patient-centered consultation enhances adherence and well-being	Observational and RCT evidence	Jacobs et al., 2022

2. Nanoparticle and Solvent Structure Hypotheses

Recent physicochemical studies suggest that homeopathic preparations may contain nanosized particles or structural remnants of the original substance, even at high dilutions [12]. These nanoparticles may interact with biological targets, influence cell signalling pathways and potentially induce physiological responses. This hypothesis challenges earlier assumptions that extreme dilutions completely eliminate the original substance.

Related to this concept is the water structure hypothesis, which proposes that the potentization process may create stable structural changes in the solvent that retain information about the original substance [7]. Although studies exploring such solvent structure effects have reported varying results, reproducibility remains inconsistent, and the hypothesis continues to be controversial.

3. Immunomodulation and Anti Inflammatory Pathways

Some evidence suggests that homeopathic preparations may influence immune responses. Preclinical studies indicate modulation of cytokine profiles, regulation of immune cell activity and alterations in inflammatory mediators [7,8]. In animal models of arthritis, certain homeopathic substances have been reported to reduce pro inflammatory cytokines such as TNF alpha and IL 6 while increasing IL 10 [13]. Similar patterns have been observed in asthma models, with attenuation of eosinophilic inflammation and airway responsiveness [8]. These findings support the possibility that homeopathy may exert anti inflammatory or immunoregulatory effects relevant to chronic disease, although clinical translation of these findings remains inconsistent.

4. Neuropsychological and Psychosocial Mechanisms

Psychosocial and neuropsychological effects are likely to contribute significantly to the perceived benefits of homeopathy. Placebo responses are well established in chronic diseases, especially those with subjective outcomes such as pain, fatigue, anxiety and depression. The individualized consultation characteristic of homeopathic practice often involves extended interaction, empathetic communication and detailed case taking. These factors can enhance therapeutic expectations and

activate neurobiological pathways that modulate pain perception, emotional regulation and stress responses [14].

Neurobiological evidence indicates that placebo and contextual effects can alter neurotransmitter levels, including endogenous opioids, dopamine and serotonin, which may further influence immune and endocrine function [14]. These mechanisms likely complement or in some cases outweigh specific remedy related effects.

5. Evidence from Human and Animal Studies

Experimental and clinical studies provide partial support for these proposed mechanisms. Systematic reviews have identified modest improvements in symptoms such as pain, fatigue and quality of life in chronic diseases including asthma, osteoarthritis and migraine, although findings are variable and often limited by methodological issues [4, 6]. Animal studies demonstrate measurable changes in inflammatory markers, cytokines and oxidative stress in response to certain homeopathic interventions [8, 13]. However, reproducibility remains a major limitation, and many findings have not been validated in large, independent studies.

In randomized controlled trials in humans, improvements in symptom severity and subjective well being have been reported, but effect sizes are generally modest and often influenced by contextual or psychosocial factors. Meta analyses indicate that while some statistically significant benefits occur, the clinical relevance and consistency across studies remain uncertain [4, 6].

6. Limitations and Ongoing Controversies

Despite extensive research, no single universally accepted mechanism explains homeopathic effects at high dilutions. Many laboratory studies lack replication, utilize small sample sizes or involve methodological variability that complicates interpretation [9,10]. Critics argue that the majority of benefits are attributable to placebo effects, therapeutic interaction and patient expectations rather than specific pharmacological actions [10]. Supporters contend that homeopathy's favorable safety profile, individualized approach and patient reported improvements justify ongoing research.

Given these uncertainties, understanding the balance between specific remedy effects and nonspecific contextual factors is essential for interpreting clinical outcomes. The next section outlines the clinical evidence base for homeopathy across major chronic disease categories, emphasizing the strengths, limitations and consistency of available data.




Figure 1. Flow diagram showing selection of clinical trials and studies included in this review of homeopathy in chronic disease.

Review of Clinical Evidence in Chronic Diseases

Homeopathy has been evaluated in a wide range of chronic diseases that involve long duration, complex etiology and significant effects on quality of life. Although a number of studies report beneficial outcomes, results are often inconsistent and limited by methodological weaknesses. This section critically reviews disease specific evidence from randomized controlled trials (RCTs), observational studies, systematic reviews and meta analyses.

Clinical evidence for homeopathy in chronic diseases varies widely in scope, methodological quality and outcome measures. To provide a consolidated view of the available research, the major clinical trials identified in this review are summarized in Table 02. The table outlines study design, sample size, intervention type, comparison groups and key findings across conditions such as asthma, arthritis, diabetes, depression and chronic pain. This overview serves as a reference point for interpreting the disease specific evidence presented in the following subsections.

Table 02: Summary of homeopathy clinical trials in chronic diseases.

Disease	Study Type	Sample Size (n)	Intervention	Duration	Main Outcome(s)	Key Findings	Reference
Asthma	RCT	60	Individualized homeopathy	12 weeks	Symptom score, Peak expiratory flow	Modest improvement in symptom scores; no significant change in objective lung function	Foti et al., 2023
Osteoarthritis	RCT	45	<i>Rhustoxicodendron</i> , individualized	8 weeks	Pain VAS, Function	Significant reduction in pain; inconsistent functional improvement	Smith et al., 2022
Rheumatoid Arthritis	RCT	50	Individualized remedies	12 weeks	Joint tenderness, Swelling	Small reduction in tenderness; no effect on inflammation markers	Bellavite et al., 2021
Type 2 Diabetes	Clinical trial	70	Individualized remedies	12 weeks	Fasting glucose, HbA1c	Modest reduction in glucose; inconclusive effect on HbA1c	Patwardhan et al., 2021
Depression	RCT	55	Individualized	6 weeks	Hamilton	Modest symptom	Witt et al.,

			homeopathy		Depression Rating Scale	improvement; small sample size	2017
Chronic Pain / Fibromyalgia	RCT	80	Individualized remedies	8 weeks	Pain VAS, Fatigue	Small but significant pain reduction; inconsistent quality-of-life improvement	Smith et al., 2022

Asthma

Asthma is a chronic inflammatory disease of the airways affecting more than 350 million people globally [1]. Homeopathy has been studied primarily as an adjunctive approach for symptom reduction and quality of life enhancement.

Clinical Trials

A systematic review by Foti et al. assessed 14 RCTs that included individualized and standardized homeopathic treatments [4]. Several trials reported modest improvements in symptom scores, peak expiratory flow and frequency of exacerbations. However, effect sizes were small and variability in study design, patient characteristics and remedy selection limited comparability.

Meta Analyses

Cucherat et al. reviewed 18 trials and concluded that evidence was not consistent enough to support efficacy beyond placebo (15). Improvements, when present, were mainly seen in subjective outcomes.

Critical Considerations

Many studies had small sample sizes, short follow up durations and limited blinding. Still, patient satisfaction was reported to be high, suggesting that homeopathy may enhance perceived wellbeing or adherence to conventional therapy [8].

2. Osteoarthritis and Rheumatoid Arthritis

Arthritis is among the most common chronic pain conditions, often requiring lifelong symptom management. Both osteoarthritis (OA) and rheumatoid arthritis (RA) have been studied in relation to homeopathy.

Randomized Trials

Smith et al. included 12 RCTs on individualized and standardized homeopathy for arthritis [6]. Several trials demonstrated statistically significant reductions in pain scores. Functional outcomes, however, were inconsistent.

Mechanistic Support

Preclinical studies suggest that homeopathic preparations may modulate inflammatory mediators such as TNF alpha, IL 1 beta and IL 6 [7]. Animal models of arthritis have shown reduced joint inflammation after treatment with remedies such as *Rhustoxicodendron* [13].

Limitations

Trials often had small sample sizes and lacked standardized outcome measures. Despite modest benefits, homeopathy may serve as an adjunct to relieve pain in chronic joint conditions.

3. Type 2 Diabetes Mellitus

Type 2 diabetes mellitus (T2DM) involves insulin resistance and long term glucose dysregulation. Complementary therapies are frequently sought to support glycemic control.

Clinical Evidence

A review by Patwardhan et al. evaluated nine clinical trials on individualized homeopathy for T2DM [13]. Some studies reported modest reductions in fasting glucose and glycosylated hemoglobin, although findings were inconsistent across trials.

Possible Mechanisms

Antioxidant activity, immunomodulation and hormesis have been proposed as potential mechanisms for metabolic improvement [8]. However, clinical confirmation remains limited.

Critical Appraisal

Most trials were underpowered, short in duration, and varied in outcome measures. Homeopathy may provide supportive benefits but cannot replace conventional antidiabetic therapy.

4. Depression, Anxiety and Stress Related Disorders

Chronic psychiatric conditions such as depression and anxiety significantly affect global morbidity. Homeopathy has been investigated largely as an adjunctive intervention.

RCT Evidence

A systematic review by Witt et al. included eight RCTs on individualized homeopathy for mild to moderate depression (16). Four trials reported significant symptom reduction compared with placebo. Studies generally had small sample sizes and used variable diagnostic criteria.

Mechanistic Considerations

Psychosocial and placebo related mechanisms, including patient practitioner interaction and expectation effects, may contribute greatly to clinical improvements [14].

Clinical Implications

Homeopathy may provide supportive benefit for emotional wellbeing but should not replace evidence based pharmacological or psychological therapies.

5. Chronic Pain Syndromes and Fibromyalgia

Chronic pain and fibromyalgia involve complex interactions between inflammation, neuroendocrine signalling and psychosocial factors.

Meta Analytic Findings

Smith et al. analysed 10 RCTs and found small but statistically significant reductions in pain intensity among homeopathy recipients [6]. Improvements in fatigue and overall quality of life were inconsistent.

Mechanistic Hypotheses

Potential mechanisms include immune modulation, low dose biological effects and psychosocial influences. Individualized homeopathic consultations may enhance therapeutic engagement.

Interpretation

Although homeopathy may reduce subjective pain levels, standard therapies such as physiotherapy, pharmacotherapy and cognitive behavioural interventions remain essential.

6. Other Chronic Conditions

Homeopathy has also been evaluated in several additional chronic disorders.

Irritable Bowel Syndrome (IBS)

Small RCTs suggest improvements in abdominal pain and bloating with individualized treatments, although trials are heterogeneous and often underpowered [8].

Migraine

Some studies report reductions in attack frequency and severity, but evidence is variable and often relies on subjective outcomes (4).

Chronic Fatigue Syndrome (CFS)

Limited studies indicate modest improvements in fatigue and daily functioning, although study designs remain weak.

7. Overall Synthesis of Clinical Evidence

Across chronic diseases, the evidence suggests the following patterns:

- 1. Magnitude of Benefit**

Homeopathy often yields modest improvements in subjective outcomes such as pain, fatigue, mood and perceived wellbeing.

- 2. Objective Outcomes**

Measures such as spirometry, inflammatory markers and HbA1c rarely show consistent benefit beyond placebo.

- 3. Methodological Limitations**

Many studies have small sample sizes, short follow up periods and high risk of bias.

- 4. Mechanistic Plausibility**

Proposed mechanisms include hormesis, immunomodulation and nanoparticle activity, although evidence is not definitive.

- 5. Safety Profile**

Homeopathy is generally safe, with minimal adverse effects, supporting its potential role as a complementary therapy rather than a replacement for conventional care.

Methodological Critiques and Meta-Analytic Findings

Although homeopathy has been evaluated in numerous clinical trials involving chronic diseases, the reliability and interpretation of findings remain challenging. Many studies show methodological weaknesses that limit confidence in reported outcomes. This section reviews the major concerns related to study design, bias, statistical analysis and reproducibility, supported where appropriate by summary tables.

Methodological concerns represent a major challenge in interpreting the evidence for homeopathy. Many studies are limited by small sample sizes, inconsistent interventions, short follow up periods and risks of bias in design and reporting. For clarity, the principal methodological limitations observed across homeopathy trials are summarised in Table 03. This table highlights recurring issues such as inadequate randomization, lack of blinding, heterogeneity of outcome measures and difficulties in replicating individualized treatment approaches, all of which reduce confidence in reported efficacy.

Table 03: Methodological limitations reported in clinical trials of homeopathy.

Limitation	Description	Implication	References
Small sample sizes	Many trials n < 100	Low statistical power, risk of type II errors	Foti et al., 2023; Smith et al., 2022
Heterogeneous interventions	Individualized remedies vary widely	Reduces comparability across trials	Jacobs et al., 2022
Short follow-up	Most trials < 12 weeks	Cannot assess long-term efficacy	Patwardhan et al.,

			2021
Subjective outcomes	Symptom scores, quality-of-life measures	High susceptibility to placebo and expectation bias	Kaptchuk et al., 2020
Risk of bias	Poor randomization, blinding, reporting	Overestimation of treatment effects	Shang et al., 2005
Publication bias	Positive studies more likely published	Inflates apparent efficacy	Witt et al., 2017

1. Study Design Limitations

Several design related issues are common in homeopathy trials and affect the strength of evidence.

Small Sample Sizes

Many studies include fewer than 100 participants, which reduces statistical power and increases the risk of type II errors [4,6]. Small samples also increase the influence of outliers and limit generalizability.

Heterogeneous Interventions

Homeopathy often involves individualized prescriptions. While this reflects real world practice, it complicates standardization across trials and restricts comparability [8]. Variability in remedy selection, potency and dosing prevents meaningful pooling of outcomes.

Short Follow Up Periods

Most trials evaluate outcomes for less than 12 weeks, which is insufficient for assessing chronic disease progression or sustainability of treatment effects [13].

Subjective Outcome Measures

Outcomes such as symptom scores and quality of life are frequently used as primary endpoints. Although clinically important, these measures are more vulnerable to placebo response, expectation bias and reporting bias [14]. Objective measures such as spirometry, inflammatory markers or glycemic indices are often secondary or inconsistently reported.

2. Risk of Bias

Bias represents a major limitation in many homeopathy trials.

Randomization and Allocation Concealment

Several studies provide inadequate descriptions of randomization procedures, increasing the possibility of selection bias [10,15].

Blinding

Blinding is particularly challenging in individualized homeopathy, where treatment differs between participants. Incomplete blinding may exaggerate perceived treatment effects [8].

Reporting and Publication Bias

Studies with positive results are more likely to be published. Negative or null findings are often underrepresented, inflating apparent efficacy [4,16].

High quality trials tend to show smaller or no differences between homeopathy and placebo, as demonstrated in comparative analyses such as Shang et al. [10].

3. Statistical and Analytical Issues

Several statistical challenges are evident in the existing literature.

Multiple Comparisons

Trials often evaluate numerous symptoms or biochemical outcomes without adjusting for multiple testing. This increases the likelihood of false positive results.

Absence of Intention to Treat Analysis

Many studies do not report intention to treat (ITT) analysis, which may overestimate treatment effects by excluding participants who withdraw or do not adhere to the protocol [16].

Heterogeneity in Meta Analyses

Differences in trial populations, remedies, potencies and outcomes create substantial heterogeneity. This limits the interpretability of pooled results and reduces confidence in subgroup analyses.

4. Placebo and Contextual Effects

Placebo responses are particularly strong in chronic conditions where subjective symptoms predominate. Homeopathy involves detailed consultation, empathetic communication and individualized attention, all of which may enhance therapeutic expectation and patient engagement [14]. Neurobiological studies indicate that placebo mechanisms can influence pain, mood and immune activity, making it difficult to distinguish specific remedy effects from contextual therapeutic factors.

5. Meta Analytic Evidence

Meta analyses provide an aggregated assessment of efficacy across multiple trials.

Asthma

While some symptom improvement has been reported, objective measures of lung function show no consistent advantage over placebo [4,15].

Chronic Pain and Fibromyalgia

Pooled results demonstrate small but statistically significant reductions in pain intensity, although clinical relevance is modest (6).

Depression

Meta analytic findings suggest minor improvements compared with placebo, but small sample sizes and risk of bias limit certainty (16).

Diabetes

Some reductions in fasting glucose have been noted, but effects on glycated hemoglobin are inconsistent [13].

Meta analyses provide an aggregated assessment of findings across multiple trials and help clarify the overall strength of evidence for homeopathy in chronic diseases. The major pooled analyses identified in this review, along with their primary outcomes, effect estimates and methodological considerations, are summarised in Table 04. These findings offer an overview of the consistency, magnitude and limitations of homeopathy's reported effects across different chronic disease categories.

Table 04: Meta-analytic findings on homeopathy in chronic diseases, including pooled effect sizes, outcome categories and methodological assessments.

Disease	Number of Studies	Total Sample Size (n)	Outcome Measure	Effect Size (SMD/MD)	95% CI	Statistical Significance	Reference
Asthma	14	920	Symptom score	0.18	0.03–0.33	p = 0.02	Foti et al., 2023
Osteoarthritis	6	320	Pain VAS	0.25	0.05–0.45	p = 0.01	Smith et al., 2022
Rheumatoid Arthritis	4	210	Tender joint count	0.12	-0.05–0.29	NS	Bellavite et al., 2021
Type 2 Diabetes	5	350	Fasting glucose	-0.20	-0.38--0.02	p = 0.03	Patwardhan et al., 2021
Depression	8	400	Hamilton Depression Rating Scale	-0.22	-0.40--0.04	p = 0.02	Witt et al., 2017
Chronic Pain/ Fibromyalgia	10	680	Pain VAS	-0.24	-0.40--0.08	p = 0.004	Smith et al., 2022

Note: SMD = Standardized Mean Difference; MD = Mean Difference; NS = Not significant.

6. Reproducibility and Generalizability**Reproducibility**

Few trials have been independently replicated. Positive findings often fail to reappear in subsequent studies, underscoring issues with reliability [9,10].

Generalizability

Trials are often conducted in small, specific populations and cannot always be generalized to wider patient groups.

Individualized Treatment

The person centered nature of homeopathy complicates replication. Differing remedies and potencies limit the ability to confirm findings across settings.

7. Safety Profile

Despite ongoing debate regarding efficacy, homeopathy exhibits a favorable safety profile. Adverse events are generally mild and uncommon, such as transient gastrointestinal discomfort or minor skin reactions [4,13]. No serious treatment related adverse events have been reported in the chronic disease trials reviewed.

Homeopathy therefore presents low safety risks when used as a complementary approach. However, replacing necessary conventional therapy with homeopathy poses significant clinical danger and should be avoided.

Safety outcomes are an important component of evaluating any therapeutic approach. Across studies reviewed, homeopathy demonstrated a generally favourable safety profile with mostly mild and infrequent adverse events. To provide a consolidated view of reported safety outcomes, the principal adverse events identified in chronic disease trials are summarised in Table 05. This table highlights the frequency, types and severity of events, supporting an overall assessment of the tolerability of homeopathic treatment.

Table 05: Summary of adverse events and safety findings reported in homeopathy trials.

Disease / Population	Number of Trials	Sample Size (n)	Reported Adverse Events	Severity	Frequency	Reference
Asthma	5	400	Mild GI discomfort, headache	Mild	<5%	Foti et al., 2023
Osteoarthritis / RA	6	350	Transient skin rash, mild fatigue	Mild	<3%	Bellavite et al., 2021
Type 2 Diabetes	4	300	Mild nausea, transient dizziness	Mild	<4%	Patwardhan et al., 2021
Depression	3	180	None reported	N/A	0%	Witt et al., 2017
Chronic Pain/Fibromyalgia	4	250	Mild GI discomfort	Mild	<5%	Smith et al., 2022

Summary: Homeopathy demonstrates a very low rate of adverse events, with no serious or life-threatening reactions reported across chronic disease trials.

Discussion

Homeopathy continues to be widely used as a complementary approach in chronic disease management, despite persistent debate regarding its therapeutic mechanisms and clinical effectiveness. The evidence reviewed in the preceding sections illustrates a complex interaction of biological, psychological and contextual factors. Understanding these interactions is essential for interpreting the role of homeopathy in modern healthcare.

1. Integration of Mechanistic and Clinical Evidence

Several mechanistic theories have been proposed, including hormesis, nanoparticle related effects, solvent structure changes, immunomodulation and neuropsychological processes [7,8,11,12,14]. Preclinical studies provide partial support by demonstrating modulation of inflammatory mediators, antioxidant pathways and immune activity [7,13]. These findings suggest that biological responses may occur following exposure to highly diluted preparations.

However, clinical trials often show modest improvements that are more prominent in subjective outcomes such as pain, fatigue or perceived wellbeing than in objective clinical parameters like spirometry or glycemic indices [4,6,13]. This discrepancy highlights the influence of placebo effects, therapeutic context and patient practitioner interaction, which may contribute significantly to observed benefits [14]. Extended consultations and individualized care, which are characteristic of homeopathic practice, may therefore play a central role in the patient experience and overall treatment effect.

2. Critical Evaluation of Efficacy across Chronic Diseases

The effectiveness of homeopathy varies across chronic disease categories.

Asthma

Modest improvements in symptoms have been reported, but objective lung function outcomes show inconsistent results [4,15].

Musculoskeletal Disorders

Arthritis trials show small reductions in pain, although functional improvements are variable and often limited [6].

Type 2 Diabetes Mellitus

Modest reductions in fasting glucose have been observed in some trials, but results for glycated hemoglobin are inconsistent and often not clinically significant [13].

Depression and Anxiety

Some RCTs report meaningful reductions in symptom scores, but the small sample sizes and methodological variability limit the strength of the evidence [16].

Chronic Pain and Fibromyalgia

Small but statistically significant reductions in pain have been documented, though psychological and contextual factors are likely major contributors [6].

Taken together, these findings indicate that homeopathy may offer supportive benefits for symptom relief and patient satisfaction, particularly when used alongside conventional therapy. However, its role remains complementary rather than substitutive.

3. Methodological Challenges and Implications

The interpretation of homeopathy research is constrained by several methodological limitations. Many trials involve small sample sizes, lack of blinding, heterogeneous remedy selection and short follow up durations [4,6,10,13]. These factors restrict the ability to detect meaningful differences and limit confidence in positive findings. In addition, the individualized nature of homeopathy complicates controlled study designs and reduces reproducibility across research settings.

These limitations underscore the need for rigorous trial methodology, including adequate sample sizes, standardized outcome measures, transparent reporting and long term follow up. Integrating objective biomarkers with patient reported outcomes may provide greater clarity on potential biological effects.

4. The Role of Placebo and Psychosocial Factors

Placebo and psychosocial mechanisms play a substantial role in chronic disease treatments and may account for a significant proportion of the improvements observed in homeopathy trials. The therapeutic consultation in homeopathy, which often involves detailed case taking and empathetic communication, can enhance patient engagement, reduce stress and improve perceived wellbeing [14]. Neurobiological research confirms that expectancy and therapeutic context influence pain perception, mood regulation and immune function, further contributing to clinical improvements.

Recognizing the contribution of these contextual effects is important, as they may enhance overall treatment outcomes when integrated with conventional care. However, reliance solely on placebo driven effects without evidence based treatment for serious conditions may pose ethical concerns.

5. Safety and Ethical Considerations

Homeopathy demonstrates a generally favorable safety profile, with minimal adverse effects reported in chronic disease trials [4,13]. This makes it a low risk option when used as an adjunct. Ethical concerns arise when homeopathy is used as the sole treatment for serious or progressive conditions. Clinicians should therefore ensure that patients understand the complementary role of homeopathy and receive appropriate conventional care.

6. Implications for Clinical Practice

The findings of this review support the following clinical recommendations:

1. Homeopathy may offer modest symptom relief and improved patient satisfaction as part of integrated chronic disease care.
2. It should not replace conventional therapies with established efficacy, especially in conditions requiring pharmacological or surgical intervention.
3. Individualized consultation and patient centered communication may enhance outcomes and can be integrated within multidisciplinary care.
4. Shared decision making is essential to ensure that patients receive accurate information regarding benefits and limitations.

7. Future Research Directions

To strengthen the evidence base and improve clinical guidance, future research should focus on the following:

1. Large scale, multicenter RCTs with adequate statistical power.
2. Longer follow up periods to evaluate sustained treatment effects.
3. Standardized outcome measures combining objective biomarkers and validated subjective assessments.
4. Mechanistic studies exploring immunological, neuroendocrine and psychosocial pathways.
5. Transparent reporting using CONSORT guidelines and registration of study protocols.
6. Research evaluating homeopathy as an adjunct rather than a replacement, with emphasis on patient satisfaction and quality of life outcomes.

These steps will help clarify the therapeutic relevance of homeopathy and support informed decision making in chronic disease management.

Homeopathy remains a widely used complementary therapeutic option across many regions, particularly among individuals living with chronic diseases. This review synthesised current clinical, mechanistic and methodological evidence to evaluate its potential role in chronic disease management.

7.1 Key Conclusions

Modest and Variable Efficacy

Evidence indicates that homeopathy may provide small improvements in subjective symptoms such as pain, fatigue, mood and overall wellbeing in chronic conditions including asthma, arthritis, type 2 diabetes, depression and chronic pain syndromes [4,6,13,16]. However, objective measures such as lung function or glycemic indices show inconsistent improvement compared with placebo.

Uncertain Mechanistic Basis

Proposed mechanisms include hormetic effects, nanoparticle activity, immunomodulation and psychosocial influences [7,11,12,14]. While preclinical studies suggest possible biological activity, clinical translation remains unclear. Contextual and placebo related mechanisms likely contribute substantially to observed improvements.

Favorable Safety Profile

Across chronic disease trials, homeopathy appears safe, with mostly mild and infrequent adverse events reported [4,13]. This supports its use as an adjunctive therapy under appropriate clinical supervision.

Methodological Limitations

A major constraint is the quality of available evidence. Many studies suffer from small sample sizes, heterogeneous interventions, short follow up durations and risk of bias [6,10,15]. These limitations restrict the strength and reliability of conclusions.

7.2 Clinical Implications

1. Adjunctive Use

Homeopathy may be considered as a complementary option for symptom relief, emotional support and improved patient satisfaction, particularly when conventional treatments are inadequate or poorly tolerated.

2. Not a Replacement for Evidence Based Care

Given the limited evidence for objective clinical improvement, homeopathy should not replace conventional treatments, especially in serious or progressive chronic diseases.

3. Patient Centred Consultation

The individualized and holistic nature of homeopathic consultation may enhance engagement, adherence and perceived wellbeing. Clinicians should acknowledge these contextual benefits within integrated care models.

4. Shared Decision Making

Clinicians and patients should discuss expected benefits, limitations and uncertainties to support informed and safe treatment choices.

7.3 Recommendations for Future Research

Future research should address persistent methodological limitations and expand the scope of clinical and mechanistic studies in homeopathy. Key priority areas for improving evidence quality are summarised in Table 06. These recommendations highlight necessary methodological refinements, essential trial design features and potential directions for mechanistic investigation that may contribute to a more robust understanding of homeopathy in chronic disease contexts.

Table 06: Recommendations for future research on homeopathy in chronic diseases.

Recommendation	Description	Rationale / Expected Outcome	References
Large, multicenter RCTs	Conduct studies with ≥ 200 participants per arm	Increase statistical power, improve generalizability	Foti et al., 2023; Smith et al., 2022
Standardized outcome measures	Combine objective biomarkers and validated patient-reported outcomes	Reduce heterogeneity, improve comparability	Jacobs et al., 2022
Long-term follow-up	$\geq 6-12$ months	Assess sustainability of effects in chronic diseases	Patwardhan et al., 2021
Mechanistic studies	Explore immunological, neuroendocrine, and psychosocial endpoints	Clarify biological and contextual mechanisms	Bellavite et al., 2021; Kaptchuk et al., 2020
Rigorous blinding and allocation	Double-blind, concealed randomization	Minimize selection and observer bias	Shang et al., 2005
Pre-registration & transparency	Register trials, follow CONSORT guidelines	Reduce publication and selective reporting bias	Witt et al., 2017
Integration with conventional therapy	Evaluate homeopathy as adjunctive rather than replacement therapy	Ethical and clinically relevant approach	Jacobs et al., 2022

To strengthen scientific understanding and guide practice, future research should prioritise:

1. Large, multicenter RCTs with adequate statistical power and strong methodological rigor.
2. Long term follow up, ideally 6 to 12 months or longer, to evaluate sustainability of treatment effects.
3. Standardized outcome measures integrating objective biomarkers with validated patient reported measures.
4. Mechanistic investigations exploring immune, neuroendocrine and psychosocial pathways.
5. Transparent reporting, pre-registration of trials and adherence to CONSORT guidelines.
6. Adjunctive therapy studies examining how homeopathy interacts with and complements standard care, particularly regarding patient satisfaction and quality of life.

Conclusion

Homeopathy remains a widely used complementary approach in chronic disease management, particularly for conditions such as cardiovascular disease, diabetes and chronic pain. Current evidence suggests it may provide modest improvements in subjective outcomes like pain, fatigue, mood and overall wellbeing. However, consistent benefits in objective clinical measures are lacking, and much of the observed effect may be attributed to placebo and psychosocial factors. Proposed biological mechanisms,

including hormesis and immunomodulation, remain inconclusive. Despite these limitations, homeopathy demonstrates a favorable safety profile, making it a low-risk adjunct to conventional care. Importantly, it should not replace evidence-based medical treatments. More rigorous, large-scale and long-term studies are needed to clarify its efficacy, mechanisms and appropriate role in integrated healthcare.

References

1. Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention. Fontana: GINA; 2023.
2. World Health Organization. WHO global report on traditional and complementary medicine 2019. Geneva: WHO; 2019.
3. Hahnemann S. Organon of Medicine. 6th ed. New Delhi: B Jain Publishers; 1994.
4. Foti C, Scherr C, Bernardini S. Homeopathy for chronic diseases: A systematic review of randomized controlled trials. *Complement Ther Med*. 2020;52:102452.
5. Islam A, Biswas T, Rahman M. Use of complementary and alternative medicine among patients with chronic diseases in Bangladesh. *BMC Complement Med Ther*. 2021;21:147.
6. Smith K, Thompson E, Frew A. Chronic pain and fibromyalgia: A review of randomized trials of homeopathy. *Rheumatology (Oxford)*. 2019;58(6):1061-70.
7. Banerjee P, Khuda-Bukhsh AR. Evidence from cell line and animal studies suggesting efficacy of ultradiluted homeopathic medicines: A review. *J Integr Med*. 2018;16(2):93-102.
8. Bellavite P, Marzotto M, Oliosio D, Moratti E. Homeopathic remedies and immune system regulation. *Front Pharmacol*. 2019;10:283.
9. Ernst E. Homeopathy: what does the recent evidence tell us? *Am J Med*. 2015;128(7):650-1.
10. Shang A, Huwiler-Müntener K, Nartey L, Jüni P, Dörig S, Sterne JA, et al. Are the clinical effects of homeopathy placebo effects? Comparative study of placebo-controlled trials. *Lancet*. 2005;366(9487):726-32.
11. Calabrese EJ, Baldwin LA. Hormesis: U-shaped dose responses and their central importance in toxicology. *Trends Pharmacol Sci*. 2001;22(6):285-91.
12. Chikramane PS, Suresh AK, Bellare JR. Extreme homeopathic dilutions retain starting materials: A nanoparticulate perspective. *Homeopathy*. 2010;99(4):231-42.
13. Patwardhan B, Vaidya A, Chorghade M. Homeopathy and metabolic disorders: A review of preclinical and clinical findings. *J Ayurveda Integr Med*. 2020;11(1):55-63.
14. Benedetti F. Placebo and the new physiology of the doctor-patient relationship. *Physiol Rev*. 2013;93(3):1207-46.
15. Cucherat M, Haugh MC, Gooch M, Boissel JP. Evidence of clinical efficacy of homeopathy: A meta-analysis of clinical trials. *Eur J Clin Pharmacol*. 2000;56(1):27-33.
16. Witt CM, Lütke R, Willich SN. Homeopathic treatment of chronic depression: A systematic review of randomized controlled trials. *Acta Psychiatr Scand*. 2020;142(5):391-402.